arXiv:hep-ph/0006348v1 30 Jun 2000

PROTON SPIN STRUCTURE AND THE LOW-ENERGY $pp \to pp\eta'$ REACTION

STEVEN D. BASS
Physik Department T39, Technische Universität München,
D-85747 Garching, Germany

Gluonic degrees of freedom induce a contact term in the effective chiral Lagrangian description of the low-energy $pp \to pp\eta'$ reaction. The strength of this contact term is, in part, related to the amount of spin carried by polarised gluons in a polarised proton.

Polarised deep inelastic scattering and η' physics provide complementary windows on dynamics induced by the axial $U_A(1)$ anomaly in QCD [1]. The flavour-singlet Goldberger-Treiman relation [2] relates the flavour-singlet axial-charge $g_A^{(0)}$ measured in polarised deep inelastic scattering to the η'–nucleon coupling constant $g_{\eta' NN}$. The large mass of the η' and the small value of $g_A^{(0)}$

$$g_A^{(0)}\Big|_{\rm pDIS} = 0.2 - 0.35 \tag{1}$$

extracted from deep inelastic scattering point to substantial violations of the OZI rule in the flavour-singlet $J^P = 1^+$ channel.

Working in the chiral limit the flavour-singlet Goldberger-Treiman relation reads

$$m g_A^{(0)} = \sqrt{\frac{3}{2}} F_0 \Bigl(g_{\eta' NN} - g_{QNN} \Bigr) \tag{2}$$

where $g_{\eta' NN}$ is the η'–nucleon coupling constant and g_{QNN} is an OZI violating coupling which measures the coupling of the topological charge density $Q = \frac{\alpha_s}{4\pi} G\tilde{G}$ to the nucleon. In Eq.(2) m is the nucleon mass and F_0 (~ 0.1GeV) renormalises the flavour-singlet decay constant. The coupling constant g_{QNN} is, in part, related [2] to the amount of spin carried by polarised gluons in a polarised proton. A large positive $g_{QNN} \sim 2.45$ is one possible explanation of the small value of $g_A^{(0)}|_{\rm pDIS}$.

It is important to look for other observables which are sensitive to g_{QNN}. OZI violation in the η'–nucleon system is a probe of the role of gluons in dynamical chiral symmetry breaking in low-energy QCD.

Working with the $U_A(1)$–extended chiral Lagrangian for low-energy QCD we find a gluon-induced contact interaction in the $pp \to pp\eta'$ reaction close to threshold[3]:

$$\mathcal{L}_{\text{contact}} = -\frac{i}{F_0^2}\, g_{QNN}\, \tilde{m}_{\eta'}^2\, \mathcal{C}\, \eta' \left(\bar{p}\gamma_5 p\right) \left(\bar{p}p\right) \tag{3}$$

Here $\tilde{m}_{\eta_0}$ is the gluonic contribution to the η' mass and $\mathcal{C}$ is a second OZI violating coupling which also features in $\eta' N$ scattering. The physical interpretation of the contact term (3) is a "short distance" (~ 0.2fm) interaction where glue is excited in the interaction region of the proton-proton collision and then evolves to become an η' in the final state. This gluonic contribution to the cross-section for $pp \to pp\eta'$ is extra to the contributions associated with meson exchange models.

What is the phenomenology of this gluonic interaction ?

Since glue is flavour-blind the contact interaction (3) has the same size in both the $pp \to pp\eta'$ and $pn \to pn\eta'$ reactions.

CELSIUS[4] have measured the ratio $R_\eta = \sigma(pn \to pn\eta)/\sigma(pp \to pp\eta)$ for quasifree η production from a deuteron target up to 100 MeV above threshold. They observed that R_η is approximately energy-independent $\simeq 6.5$ over the whole energy range. The value of this ratio signifies a strong isovector exchange contribution to the η production mechanism[4]. This experiment should be repeated for η' production. The cross-section for $pp \to pp\eta'$ close to threshold has been measured at COSY[5]. The $pn \to pn\eta'$ measurement remains to be done. The more important that the gluon-induced process (3) is in the $pp \to pp\eta'$ reaction the more one would expect $R_{\eta'} = \sigma(pn \to pn\eta')/\sigma(pp \to pp\eta')$ to approach unity near threshold after we correct for the final state interaction between the two outgoing nucleons. (After we turn on the quark masses, the small $\eta - \eta'$ mixing angle $\theta \simeq -18$ degrees means that the gluonic effect (3) should be considerably bigger in η' production than η production.)

η' phenomenology is characterised by large OZI violations. It is natural to expect large gluonic effects in the $pp \to pp\eta'$ process. A measurement of $R_{\eta'}$ is urgently needed!

References

1. S.D. Bass, hep-ph/9902280, Eur. Phys. J **A5** (1999) 17.
2. G. Veneziano, Mod. Phys. Lett. **A4** (1989) 1605;
   G.M. Shore and G. Veneziano, Nucl. Phys. **B381** (1992) 23.
3. S.D. Bass, hep-ph/9907373, Phys. Lett. **B463** (1999) 286.
4. The CELSIUS Collaboration (H. Calen et al.), Phys. Rev. **C58** (1998) 2667.
5. The COSY-11 Collaboration (P. Moskal et al.), Phys. Rev. Lett. **80** (1998) 3202; Phys. Lett. **B474** (2000) 416.